




CytoGrow™ Recombinant R-Spondin1 (Human)

Catalog Numbers: CG067-A, CG067-B, CG067-C Revision 1.0

Specifications

luman 0µg/50µg/1mg ag Free 95%, by SDS-PAGE (under reducing (R) & Non-reducing conditions, visualized by Coomassie staining) 1EU/mg by the LAL method HO er21-Ala263 .0-8.0
ag Free 95%, by SDS-PAGE (under reducing (R) & Non-reducing conditions, visualized by Coomassie staining) 1EU/mg by the LAL method HO er21-Ala263
95%, by SDS-PAGE (under reducing (R) & Non-reducing conditions, visualized by Coomassie staining) 1EU/mg by the LAL method HO er21-Ala263
1EU/mg by the LAL method HO er21-Ala263
H0 er21-Ala263
er21-Ala263
.0-8.0
hite powder, Colorless clear liquid after reconstitution
.22 μm-filtered solution containing PBS, pH 7.4
he product is shipped with blue ice.
is recommended to redissolve in sterile deionized water.
yophilized state: 6 months at -20°C to -80°C fter reconstitution and under sterile conditions. months at -20°C to -80°C
6



Biological activity is assessed based on the induction of TopFlash reporter activity in CHO cells in the presence of 50 ng/mL Wnt3a. The expected ED₅₀ is less than 1.0 µg/mL.

Reconstitution

- Reconstitute protein to concentrations of 100 ug/mL or higher.
- Use sterile deionized water to solubilize the lyophilized protein.
- Gently tap glass vial to help collect the powder to the bottom.
- Add desired amount of buffer, observe after 2 min. Gentle agitation by hand may be required.
- Prepare aliquots of solution. Freeze at -20°C to -80°C to avoid repeated freeze thaw cycles.